

TABLE
Number of mosquito species collected by larval habitat, size, grass, substratum and water characteristics in immature stages habitats sampled in the province of Chaco, Argentina

Group	Species	Larval habitat								Water						
		Bromeliad		Lagoon	Pool	Ditch	Deep pool	Artificial container	Size (m)	Grass	Substrate	Temperature °C (mean)	Colour	pH (mean)	Permanence	Depth
Terrestrial	Epiphytic															
1	<i>Aedes aegypti</i>	-	-	-	-	-	-	466	0.5-0.65 x 0.65-1.5	A	-	19-37 (29.75)	B/R	7.6-8.12 (7.93)	TE	0.01-0.03
2	<i>Culex imitator</i>	2,281	206	-	-	-	-	-	0.9-1.5	A	-	6-39 (25.76)	T/Y/R	4-8.50 (6.24)	P	0.01-0.25
	<i>Wyeomyia muehlensi</i>	214	9	-	-	-	-	-	1.2-1.5	A	-	6-32 (23.56)	T/Y/R	4-8.5 (5.98)	P	0.01-0.15
	<i>Toxorhynchites haemorrhoidalis separatus</i>	74	6	-	-	-	-	-	0.9-1.5	A	-	16-39 (25.21)	T/Y/R	4-7.4 (5.83)	P	0.02-0.08
	<i>Culex davisi</i>	49	9	-	-	-	-	-	0.9-1.5	A	-	6-33 (24.90)	T/Y/R	4-7.0 (6.02)	P	0.01-0.06
3A	<i>Psorophora ciliata</i>	-	-	-	113	6	1	-	0.1-3 x 0.15-50	PR/A	M	20-38 (27.67)	B	6-8.1 (6.60)	TE/SP	0.01-0.23
	<i>Psorophora varinervis</i>	-	-	-	330	-	-	-	0.15-7 x 0.15-35	PR/A	M	22-40 (31.16)	B	6-8.28 (7.36)	TE/SP	0.05-0.35
	<i>Psorophora confinnis</i>	-	-	-	460	1	32	-	0.08-5 x 0.09-35	PR/A	M	7-40 (29.46)	B	6-8.83 (7.30)	TE/SP	0.01-0.64
	<i>Psorophora cyanescens</i>	-	-	-	377	15	8	-	0.1-8 x 0.15-35	PR/A	M	20-35 (28.25)	T/B	6-8.44 (6.95)	TE/SP	0.04-0.23
	<i>Psorophora cingulata</i>	-	-	-	491	9	86	-	0.15-50	PR/A	M	7-40 (27.31)	T/B	6-8.58 (7.40)	TE/SP	0.02-0.27
	<i>Psorophora ferox</i>	-	-	-	364	-	-	-	0.1-8 x 0.12-35	PR	M	10-34 (25.54)	B	5-8.11 (7.14)	TE/SP	0.01-0.25
	<i>Ochlerotatus hastatus/oligopistus</i>	-	-	-	875	-	-	-	0.1-4.5 x 0.1-50	PR/A	M	10-35 (24.47)	T/B	6-8.28 (7.02)	TE/SP	0.03-0.27
	<i>Ochlerotatus serratus</i>	-	-	-	635	5	41	-	0.12-5 x 0.12-30	PR/A	M	9-34 (24.6)	T/B	6-8.21 (7.15)	TE/SP	0.04-0.20
	<i>Ochlerotatus scapularis</i>	-	-	-	578	68	4	-	0.12-10 x 0.12-30	PR/A	M/S	7-32 (22.2)	T/B	6-8.58 (7.11)	TE/SP	0.02-0.27
	<i>Culex intricatus</i>	-	-	3	82	2	-	-	0.10-4 x 0.1-25	PR/A	M	15-34 (23.33)	B	5-8.60 (7.34)	TE/SP/P	0.04-0.25
	<i>Culex quinquefasciatus</i>	-	-	-	11	208	1,350	71	0.2-10 x 0.9-30	PR	M	9-33 (23.12)	B	6-8.50 (8.23)	TE/SP/P	0.02-0.18
	<i>Culex pilosus</i>	-	-	12	334	6	3	-	0.15-10 x 0.15-30	PR/A	M	14-35 (25.91)	T/B	5-8.58 (7.07)	TE/SP/P	0.02-0.25
	<i>Ochlerotatus albifasciatus</i>	-	-	-	185	89	-	-	0.2-10 x 0.25-30	PR/A	M/S	12-32 (21.32)	T/B	6-8.30 (7.19)	TE/SP	0.02-0.25
	<i>Culex bidens</i>	-	-	27	158	248	4	-	0.3-10 x 0.1-50	PR	M/S	7-38 (22.99)	T/B	5-8.67 (7.26)	TE/SP/P	0.03-0.28
3B	<i>Culex maxi</i>	-	-	241	1,343	759	261	-	0.1-10 x 0.1-40	PR/A	M/S	7-40 (23.48)	T/B	5-8.74 (6.80)	TE/SP/P	0.01-0.28
	<i>Uranotaenia lowii</i>	-	-	65	91	317	-	-	0.1-10 x 0.1-50	PR	S	9-36 (21.72)	T/B	6-8.61 (7.45)	TE/SP/P	0.02-0.27
	<i>Culex eduardoi</i>	-	-	34	49	190	-	-	0.1-15 x 0.1-35	PR	S	7-34 (19.07)	T/B	6-8.28 (7.27)	SP/P	0.04-0.80
	<i>Culex chidesteri</i>	-	-	37	22	22	2	-	0.1-15 x 0.1-50	PR	M/S	12-34 (22.32)	T/B	6-8.98 (7.45)	SP/P	0.04-0.28
	<i>Anopheles neomaculipalpus</i>	-	-	9	85	-	-	-	0.1-8 x 0.1-50	PR/A	M	9-39 (25.50)	B	5-8.15 (7.02)	TE/P	0.02-0.10
	<i>Uranotaenia pulcherrima</i>	-	-	18	17	-	-	-	0.10-5 x 0.10-50	PR	M/S	8-32 (25.06)	B	6-8.43 (6.89)	TE/P	0.03-0.25
	<i>Anopheles triannulatus</i>	-	-	218	110	7	-	-	0.10-6 x 0.10-50	PR	M/S	8-40 (24.11)	T/B	6-8.77 (7.31)	TE/SP/P	0.05-0.37
	<i>Anopheles albitarsis</i>	-	-	69	129	4	-	-	0.10-6 x 0.10-50	PR/A	M/S	7-39 (23.52)	T/B	6-8.98 (7.19)	TE/SP/P	0.10-0.35
	<i>Uranotaenia apicalis</i>	-	-	86	2	3	1	-	0.10-5 x 0.10-50	PR	S	20-36 (28.65)	B	6-8.22 (7.37)	SP/P	0.03-0.27
	<i>Mansonia humeralis</i>	-	-	59	-	-	-	-	3.88-4 x 30-50	A	S	15-40 (27.38)	B	5-7.80 (7.21)	P	0.09-0.32
	<i>Aedeomyia squamipennis</i>	-	-	105	-	-	-	-	3.88-4 x 30-50	A	S	18-40 (28.25)	B	6-7.97 (7.32)	P	0.09-0.30

A: absence; B: brown; M: muddy; P: permanent; PR: presence; R: reddish; S: sandy; SP: semi-permanent; T: transparent; TE: temporary; Y: yellowish.