

Levels of IgG and its subclasses measure in patients with paracoccidioidomycosis (PCM) compared according to their scheme of treatment

Condition of treatment	Mexo (mean ± SD)					rPb27 (mean ± SD)				
	Total IgG	IgG1	IgG2	IgG3	IgG4	Total IgG	IgG1	IgG2	IgG3	IgG4
SMZ-TMP	1.596 ± 1.280	0.308 ± 0.322	0.397 ± 0.477	0.175 ± 0.131	0.145 ± 0.168	0.821 ± 0.764	0.502 ± 0.823	0.291 ± 0.491	0.122 ± 0.094	0.311 ± 0.655
Ceto	0.919 ± 0.556	0.167 ± 0.152	0.217 ± 0.200	0.106 ± 0.044	0.115 ± 0.057	0.617 ± 0.512	0.293 ± 0.418	0.151 ± 0.089	0.102 ± 0.042	0.181 ± 0.241
Itra	0.919 ± 0.556	0.254 ± 0.152	0.725 ± 0.523	0.269 ± 0.294	0.180 ± 0.165	0.749 ± 0.520	0.328 ± 0.302	0.218 ± 0.092	0.110 ± 0.061	0.255 ± 0.450
NT	0.919 ± 0.556	0.881 ± 0.997	0.799 ± 0.898	0.205 ± 0.196	0.323 ± 0.284	1.250 ± 1.073	0.527 ± 0.915	0.158 ± 0.061	0.141 ± 0.087	0.564 ± 0.912

ceto: cetoconazole; itra: itraconazole; mean ± standard deviation (SD): median of levels of IgG and its subclasses plus its SD; NT: not treated patients; rPb27: recombinant Pb27; SMZ-TMP: sulphamethoxazole-trimethoprim.

Levels of IgG and its subclasses measured in patients with paracoccidioidomycosis (PCM) compared according to their clinical manifestations

Clinical form	Mexo (mean ± SD)					rPb27 (mean ± SD)				
	Total IgG	IgG1	IgG2	IgG3	IgG4	Total IgG	IgG1	IgG2	IgG3	IgG4
Unifocal	1.254 ± 0.735	0.364 ± 0.576	0.537 ± 0.562	0.145 ± 0.118	0.206 ± 0.223	0.821 ± 0.734	0.525 ± 0.868	0.338 ± 0.609	0.146 ± 0.117	0.271 ± 0.613
Multifocal	1.154 ± 0.789	0.308 ± 0.449	0.361 ± 0.412	0.181 ± 0.142	0.127 ± 0.132	0.717 ± 0.659	0.398 ± 0.612	0.143 ± 0.094	0.100 ± 0.048	0.278 ± 0.558

mean ± standard deviation (SD): median of levels of IgG and its subclasses plus its SD; rPb27: recombinant Pb27.