

## SHORT COMMUNICATION

## First report of *Lymnaea cousini* Jousseau, 1887 naturally infected with *Fasciola hepatica* (Linnaeus, 1758) (Trematoda: Digenea) in Machachi, Ecuador

Ángel Villavicencio A/<sup>+</sup>, Mauricio Carvalho de Vasconcellos \*

Department of Veterinary Pathology, Agrary Faculty, Russian People's Friendship University, Moscow-Russia, 117-198, St. M. Maklaya 6 \*Núcleo de Biologia e Controle de Endo e Ectoparasitas de Interesse Médico e Veterinário, Departamento de Biologia, Instituto Oswaldo Cruz-Fiocruz, Rio de Janeiro, RJ, Brasil

*We report the first finding of Lymnaea cousini naturally infected with Fasciola hepatica in Ecuador. A sample of 70 snails was collected in April 2005 from a wetland located in a valley at approximately 3000 m a.s.l., near the locality of Machachi, Pichincha Province. The prevalence of natural infection in L. cousini was 31.43%, which is the highest value ever recorded for naturally infected lymnaeid species.*

Key words: *Lymnaea cousini* - *Fasciola hepatica* - Ecuador

Fascioliasis, one of the main diseases of ruminants, is caused by the liver fluke *Fasciola hepatica*. This worldwide distributed disease occasionally affects humans and produces an important economic impact on livestock due to reduced weight gain, lowered fertility and abortion, progressive decrease in milk yield, as well as liver condemnation. The annual losses to the agricultural sector are estimated to be US\$2.2 million, with over 600 million animals infected worldwide (Dalton 1999).

In Ecuador the present prevalence of *F. hepatica* in livestock remains uncertain because most data are collected by sanitary inspectors from slaughter houses. The Andean region is the endemic area of fascioliasis, where prevalence of infection in livestock ranges from 20 to 60% and in humans from 24 to 53% (SESA 2003). This region is inhabited by 23.6 and 52.9% of the total and rural population, respectively; almost 200,000 people are infected and 1% are at risk of infection (WHO 1995).

Despite fascioliasis being a serious health problem in the country, very little is known about snails of the genus *Lymnaea*, the intermediate hosts responsible for *F. hepatica* transmission.

Some of the lymnaeid species occurring in Andean countries are as follows: *L. truncatula* Müller, 1774 from Bolivian highlands (Jabbour-Zahab et al. 1997, Mas-Coma et al. 1999) and Venezuela (Pérez Mata 2003); *L. columella* Say, 1817 from Venezuela and Ecuador (Pérez Mata 2003); *L. cubensis* Peiffer, 1839 from Venezuela (Pérez Mata 2003) and *L. cousini* Jousseau, 1887 (syn. *L. ubaquensis*

Piaget, 1912; *L. bogotensis* Pilsbry, 1935; see Hubendick 1951 and Pointier et al. 2004). *L. cousini* has been found in Ecuador (Jousseau 1887, Paraense 1995, 2004), Colombia (Piaget 1912, Pilsbry 1935), and Venezuela (Pointier et al. 2004), but no report concerning natural infection has ever been published.

The aim of this study is to report the first finding of *L. cousini* naturally infected with *F. hepatica* in Ecuador.

A sample of 70 snails was collected in April 2005 within a private farm where fascioliasis in cattle was previously diagnosed by faecal examination. It was located near Machachi (78°30'W, 00°30'S, 3100 m a.s.l.) in the Pichincha Province. The sampling site was a wetland placed in a relatively low valley at approximately 3000 m a.s.l. and covered an area of 4.5 m<sup>2</sup>. Water in the wetland flowed slowly, had a maximum height of 10 cm and pH was 6.5. The soil is volcanic in origin, and vegetation was composed of Ray-grass (*Lolium* sp.) and aquatic plants.

Snails collected were kept alive in plastic containers with wet cotton pads and transported to the Department of Malacology of the Instituto Oswaldo Cruz, Rio de Janeiro, Brazil. Upon arrival at the laboratory, ten specimens were relaxed, killed, and preserved according to Paraense (1984), and their taxonomic identification was performed by Dr Paraense on the basis of features of the shell and internal organs (Paraense 1995). The remaining snails shared the same external appearance as those already determined. Specimens were examined for the presence of larval stages of *F. hepatica* under stereoscopic microscope.

All 70 collected snails belonged to *L. cousini* Jousseau, 1887 (Figs 1, 2), and 22 (31.43%) were infected with *F. hepatica* (Fig. 3).

Shells of *L. cousini* specimens from Ecuador described by Jousseau (1887) ranged 10-14 mm in height and 6-10 mm in diameter with an aperture of 7-10 mm long, while those from Venezuela varied 6.6-9.3 mm in height and 3.7-6 mm in width (Pointier et al. 2004).

Financial support: Fundacyt, Proj. 2005701

<sup>+</sup>Corresponding author. Email: fveterinario@hotmail.com

Received 22 July 2005

Accepted 7 November 2005

The surprisingly high prevalence of natural infection with *F. hepatica* in *L. cousini* may be explained by environmental conditions favouring the development of snail populations and the transmission of the parasite. Valleys in the Andean region of Ecuador are characterized by tem-



Fig. 1: *Lymnaea cousini* from Machachi, Ecuador.



Fig. 2: shells of *Lymnaea cousini* collected in Machachi, Ecuador.



Fig. 3: redia and cercaria of *Fasciola hepatica* in naturally infected *Lymnaea cousini* from Machachi, Ecuador. 50×

peratures ranging from 10 to 30°C, frequent precipitation, numerous streams, and low-permeable soil layers. In addition, anthropogenic factors involving irrigation of pasture lands by flood or ditches and inappropriate management of grazing cattle are unlikely to prevent infection in both intermediate and definite hosts.

The prevalence of *F. hepatica* in *L. cousini* from Machachi, Ecuador (31.43%), is the highest value ever reported for lymnaeid snails naturally infected with this parasite. In Brazil, *L. columella* showed the following infection rates: 2.4% in Rio de Janeiro (Rezende et al. 1973); 5.2 and 3.9% in Minas Gerais (Coelho & Lima 2003); 1.22 and 0.14% (Ueta 1980), 5.26% (Oliveira et al. 2002) in São Paulo; 3.3% in Rio Grande do Sul (Silva Santos et al. 1987). In Corrientes, Argentina, the prevalence of *L. columella* was 8.8% (Prepelitchi et al. 2003). In central France, Rondelaud et al. (2001) reported a global prevalence of natural infection of 1.1% in *L. truncatula* and 0.3% in *L. glabra*. In the Russian Federation, *L. truncatula* populations showed infection prevalences ranging from 0.05 to 0.72% (Villavicencio 2004).

#### ACKNOWLEDGEMENTS

To Dr Wladimir Lobato Paraense, Department of Malacology, Instituto Oswaldo Cruz-Fiocruz, for his valuable help in identifying the snails. To Natalia Dondua for field assistance. To AF Barbosa for the photographs, and to all the people who lent their support and assistance during this study.

#### REFERENCES

- Coelho LHL, Lima WS 2003. Population dynamics of *Lymnaea columella* and its natural infection by *Fasciola hepatica* in the state of Minas Gerais, Brazil. *J Helminthol* 77: 7-10.
- Dalton JP 1999. *Fasciolosis*, CABI publishing, New York, 562 pp.
- Hubendick B 1951. Recent Lymnaeidae. Their variation, morphology, taxonomy, nomenclature, and distribution. *Kungl Svenska Vetenskapsakad Handl* (4 ser) 3: 1-223.
- Jabbour-Zahab R, Pointier JP, Jourdane J, Jarne P, Oviedo JA, Bargues MD, Mas-Coma S, Anglés R, Perera G, Balzan C, Khallayoune K, Renaud F 1997. Phylogeography and genetic divergence of some Lymnaeid snails, intermediate hosts of human and animal fascioliasis with special reference to lymnaeids from the Bolivian Altiplano. *Acta Trop* 64: 191-203.
- Jousseume F 1887. Mollusques nouveaux de la République de l'Equateur. *Bull Soc Zool France* 12: 165-186.
- Mas-Coma S, Anglés R, Esteban JG, Bargues MD, Buchon P, Franken M, Strauss W 1999. The Northern Bolivian Altiplano: a region highly endemic for human fascioliasis. *Trop Med Int Hlth* 4: 454-467.
- Oliveira SM, Fujii TU, Sposito Filha E, Martins A 2002. Ocorrência de *Lymnaea columella* Say, 1817 infectada naturalmente por *Fasciola hepatica* (Linnaeus, 1758), no Vale do Ribeira, São Paulo, Brasil. *Arq Inst Biológico* 69: 29-37.
- Paraense WL 1984. *Lymnaea diaphana*: a study of topotypic specimens (Pulmonata: Lymnaeidae). *Mem Inst Oswaldo Cruz* 79: 75-81.

- Paraense WL 1995. *Lymnaea cousini* Jousseume, 1887, from Ecuador (Gastropoda: Lymnaeidae). *Mem Inst Oswaldo Cruz* 90: 605-609.
- Paraense WL 2004. Planorbidae, Lymnaeidae and Physidae of Ecuador (Mollusca: Basommatophora). *Mem Inst Oswaldo Cruz* 99: 357-362.
- Pérez Mata AM 2003. *Contribuciones a la Caracterización de la Fascioliasis en Países Andinos, Transmisión, Parásitos y Vectores en Venezuela, Perú y Ecuador*. PhD Thesis, Facultad de Farmacia, Universidad de Valencia, España, 544 pp.
- Piaget J 1912. Quelques mollusques de Colombie. *Mém Soc Neuchatel Sci Nat* 5: 265-269.
- Pislbry HA 1935. South American land and freshwater molluscs, IX-Colombian species. *Proc Acad Nat Sci Philadelphia* 87: 83-88.
- Pointier JP, Noya O, Amarista M, Théron A 2004. *Lymnaea cousini* Jousseume, 1887 (Gastropoda: Lymnaeidae): first record for Venezuela. *Mem Inst Oswaldo Cruz* 99: 567-569.
- Prepelitchi L, Florencia Kleiman, Silvia M Petrokovsky, Ricardo A Moriena, Oscar Racioppi, José Alvarez, Cristina Wisnivesky-Colli 2003. First report of *Lymnaea columella* Say, 1817 (Pulmonata: Lymnaeidae) naturally infected with *Fasciola hepatica* (Linnaeus, 1758) (Trematoda: Digenea) in Argentina. *Mem Inst Oswaldo Cruz* 98: 889-891.
- Rezende HEB, Araújo JL, Gomes PAC, Nurenberg S, Neto MP, Oliveira GP, Mello RP 1973. Notas sobre duas espécies de *Lymnaea* Lamark, 1799, hospedeiros intermediários de *Fasciola hepatica* L. no estado do Rio de Janeiro (Mollusca, Gastropoda, Basommatophora, Lymnaeidae). *Arq Univ Fed Rural* 3: 21-23.
- Rondelaud D, Vignoles P, Abrous M, Dreyfuss G 2001. The definitive and intermediate hosts of *Fasciola hepatica* in the natural watercress beds in central France. *Parasitol Res* 87: 475-478.
- Silva Santos S, Laranja RJ, Costa NC, Bothelo GA, Ceresér VH, Martins JR 1987. *Lymnaea columella* Say, 1817 interagindo na disseminação de fasciolose na Estação Experimental Zootécnica de Dom Pedrito, RS. *Hora Veterinária* 6: 42-48.
- SESA - Servicio Ecuatoriano de Sanidad Animal 2003. *Estadísticas Anales de Control Epidemiológico*, Quito, 90 pp.
- Ueta MT 1980. Ocorrência de infecção natural de *Fasciola hepatica* Linnaeus, 1758 em *Lymnaea columella* Say, 1817 no Vale do Paraíba, SP, Brasil. *Rev Saúde Pública* 14: 230-233.
- Villavicencio A 2004. Infección natural de moluscos *L. truncatula* con estados larvários de *Fasciola hepatica* en regiones de Moscú. *Material Conf Cient sobre la Lucha contra las Zoonosis* 5: 199-122.
- WHO-World Health Organization 1995. Control of foodborne trematode infections. Report of a WHO Expert Committee. WHO Technical Report Series, Geneva No. 84, 200 pp.