

# Decreased memory T-cell response and function in human immunodeficiency virus-infected patients with tegumentary leishmaniasis

Luana Leandro Gois<sup>1</sup>, Sanjay Mehta<sup>2</sup>, Robert T Schooley<sup>2</sup>,  
Roberto Badaró<sup>2,3</sup>, Maria Fernanda Rios Grassi<sup>1/+</sup>

<sup>1</sup>Centro de Pesquisa Gonçalo Moniz-Fiocruz, Salvador, BA, Brasil <sup>2</sup>University of California San Diego, La Jolla, CA, USA

<sup>3</sup>Universidade Federal da Bahia, Salvador, BA, Brasil

*The effects of human immunodeficiency virus (HIV) on the immune response in patients with cutaneous leishmaniasis have not yet been fully delineated. This study quantified and evaluated the function of memory T-cell subsets in response to soluble Leishmania antigens (SLA) from patients coinfecting with HIV and Leishmania with tegumentary leishmaniasis (TL). Eight TL/HIV coinfecting subjects and 10 HIV seronegative subjects with TL were evaluated. The proliferative response of CD4<sup>+</sup> and CD8<sup>+</sup> T-cells and naive, central memory (CM) and effector memory (EM) CD4<sup>+</sup> T-cells in response to SLA were quantified using flow cytometry. The median cell division indices for CD4<sup>+</sup> and CD8<sup>+</sup> T-cells of coinfecting patients in response to SLA were significantly lower than those in patients with Leishmania monoinfection ( $p < 0.05$ ). The proportions of CM and EM CD4<sup>+</sup> T-cells in response to SLA were similar between the coinfecting patients and patients with Leishmania monoinfection. However, the median CM and EM CD4<sup>+</sup> T-cell counts from coinfecting patients were significantly lower ( $p < 0.05$ ). The reduction in the lymphoproliferative response to Leishmania antigens coincides with the decrease in the absolute numbers of both EM and CM CD4<sup>+</sup> T-cells in response to Leishmania antigens in patients coinfecting with HIV/Leishmania.*

Key words: *Leishmania* - HIV - coinfection - memory CD4<sup>+</sup> T-cells

Human immunodeficiency virus (HIV) type 1 is a human retrovirus that infects roughly 34 million people worldwide and is responsible for the majority of the acquired immune deficiency syndrome (AIDS) epidemic. The most severely affected areas are in Sub-Saharan Africa, the Caribbean, Eastern Europe and Central Asia (UNAIDS 2012). *Leishmania* have been recognised as opportunistic pathogens in immunocompromised individuals, including those infected with HIV (Badaro 1997, Cruz et al. 2006). Over the past decades, the HIV epidemic has spread from urban areas to suburban and rural areas, where leishmaniasis is found. In particular, *Leishmania* and HIV infection overlap in several subtropical and tropical regions around the world (Karp & Auwaerter 2007). In northern India, Sudan and Ethiopia, visceral leishmaniasis has been a major scourge in patients infected with HIV (Wolday et al. 2001, Mathur et al. 2006, ter Horst et al. 2008). In Brazil, cutaneous disease is also a common presentation of leishmaniasis in HIV-coinfecting patients (Rabello et al. 2003, Carranza-Tamayo et al. 2009, Lindoso et al. 2009). These patients often present with multiple skin ulcers, disseminated le-

sions and nasopharyngeal and/or genital mucosal damage (Coura et al. 1987, Machado et al. 1992, Da-Cruz et al. 1999, Lindoso et al. 2009). The clinical outcome of *Leishmania* infection is dependent upon the balance of T-helper (Th)1 and Th2 cytokine production during the immune response (Bacellar et al. 2000, Bourreau et al. 2003). In HIV/*Leishmania* coinfecting patients with visceral disease, HIV infection alters the Th1/Th2 balance, resulting in an increased Th2 cytokine response while levels of the Th1 cytokines interleukin (IL)-12 and IL-18 are decreased (Nigro et al. 1999, Wolday et al. 2000). This results in more aggressive and relapsing visceral leishmaniasis. The effects of HIV on the immune response in cutaneous leishmaniasis have not yet been fully delineated. Reduced interferon (IFN)- $\gamma$  production, decreased IFN- $\gamma$ /IL-10 ratios (Rodrigues et al. 2011) and diminished lymphoproliferative responses to *Leishmania* antigens have been described in HIV-infected patients with the cutaneous form of leishmaniasis (Da-Cruz et al. 1992, 1999). These findings suggest that coinfecting HIV/*Leishmania* patients may have a memory T-cell defect. Here, we have quantified and assessed the function of memory T-cell subsets in response to soluble *Leishmania* antigens (SLA) from patients coinfecting with HIV and *Leishmania* with tegumentary leishmaniasis (TL).

## SUBJECTS, MATERIALS AND METHODS

Eight patients with TL and HIV infection and 10 HIV seronegative subjects with TL were included in the study. The subjects were enrolled from the AIDS clinical centre of the University Hospital Professor Edgard Santos, located in Salvador, Brazil, after informed consent.

doi: 10.1590/0074-0276130174

Financial support: UCSD Center for AIDS Research, NIH (P30AI036214)

+ Corresponding author: grassi@bahia.fiocruz.br

Received 27 March 2013

Accepted 18 July 2013

Clinical and demographic characteristics of the HIV/*Leishmania* coinfecting subjects are presented in Table I. All *Leishmania* monoinfected patients had active cutaneous leishmaniasis at the time of inclusion in the study. Six of these patients had cutaneous leishmaniasis, one had disseminated cutaneous leishmaniasis and three had mucosal leishmaniasis.

Peripheral blood mononuclear cells (PBMCs) were isolated by passage over a Ficoll-Hypaque gradient (Amersham Biosciences, Piscataway, NJ, USA). PBMCs ( $3 \times 10^6$  cells) were resuspended in RPMI-1640 medium (Sigma-Aldrich, St. Louis, MO, USA) supplemented with 2 mM L-glutamine, penicillin (100 U/mL), streptomycin (100 mg/mL) (GIBCO) and 5% heat-inactivated human AB serum. The cells were stimulated with SLA obtained from the *Leishmania chagasi* (MHOM/BR2000/Merivaldo2) strain [isolated from a patient with visceral leishmaniasis (Baleeiro et al. 2006)] or phytohemagglutinin (PHA) (Sigma-Aldrich). Control cultures were also established with medium alone.

To evaluate the proliferative response of T-cell subsets to SLA,  $1.5 \times 10^6$  PBMC/mL were labelled with carboxyfluorescein succinimidyl ester (Invitrogen, Eugene, OR, USA) as described by the manufacturer. PBMCs were cultured in  $12 \times 75$ -mm tissue culture tubes for five days at 37°C in a humidified 5% CO<sub>2</sub> atmosphere. Then, the cells were washed with phosphate-buffered saline (PBS)/bovine serum albumin and surface stained with anti-CD4-phycoerythrin (PE) (Becton Dickinson, CA, USA) and anti-CD8-PE-Cy5 (eBioscience, CA, USA) for 30 min at 4°C. After fixation in 1% formaldehyde, the cells were acquired by FACS Aria (Becton Dickinson) and analysis was performed using FlowJo software version 7.6 (Tree Star Inc, Ashland, OR, USA). At least 30,000 events were analysed per sample. The proliferation intensity was determined through the percentage of the cell division index. The threshold to define positive proliferative response was a cell division index above 0.06 for CD4<sup>+</sup> T-cells and above 0.09 for the CD8<sup>+</sup> T-cell subset as described previously (Pinto et al. 2011).

Alternatively, to determine the frequency of memory CD4<sup>+</sup> T-cells,  $1.5 \times 10^6$  PBMCs were cultured with SLA, PHA or PBS for 18 h in supplemented RPMI-1640 medium in the presence of 0.5 µg of purified anti-CD28 antibody at 37°C in a humidified 5% CO<sub>2</sub> atmosphere. Then, the cells were stained with anti-CD4-(fluorescein isothiocyanate), anti-CD45RA-PE, anti-CD62L-PE-Cy5 and anti-CCR7-PE-Cy7 antibodies (Becton Dickinson). Cells were acquired by FACS Aria and analysed using the FlowJo software. Naïve cells were defined as CD4<sup>+</sup>CD45RA<sup>+</sup>CD62L<sup>+</sup>CCR7<sup>+</sup>, central memory (CM) cells were defined as CD4<sup>+</sup>CD45RA<sup>-</sup>CD62L<sup>+</sup>CCR7<sup>+</sup> and effector memory (EM) cells were defined as CD4<sup>+</sup>CD45RA<sup>-</sup>CD62L<sup>-</sup>CCR7<sup>-</sup>. Absolute naïve, CM and EM lymphocyte subpopulation counts were obtained by multiplying the subpopulation percentage by the absolute lymphocyte count.

The data are expressed as median and interquartile range (IQR). Statistical analysis was performed using GraphPad Prism 5.02 (La Jolla, CA, USA). The Mann-Whitney *U* test was used for group comparisons and the Wilcoxon test was used for the analyses of cell count and cell proportion in the presence and absence of antigen.

Institutional review board approvals were obtained from Brazilian National Ethical Commission, Oswaldo Cruz Foundation and the University of California San Diego Human Research Protection Program.

## RESULTS

The median CD4<sup>+</sup> T-cell count from HIV/*Leishmania* coinfecting patients (241 cells/mm<sup>3</sup>, IQR 85-453) was significantly lower compared with that of *Leishmania* monoinfected patients (784 cells/mm<sup>3</sup>, IQR 599-1477) ( $p = 0.004$ ). The median HIV viral load of HIV/*Leishmania* coinfecting patients was 4.6 log copies/mL (IQR 1.9-5.2). The sex ratio and median age were similar in both groups.

The median cell division indices in response to SLA for CD4<sup>+</sup> (0.06, IQR 0.05-0.1) and CD8<sup>+</sup> T-cells (0.05, IQR 0.01-0.08) from HIV/*Leishmania* coinfecting patients were significantly lower than those from *Leishmania* monoinfected patients (CD4<sup>+</sup> T-cell: 0.18, IQR 0.15-0.26;

TABLE I

Clinical and epidemiological characteristics of human immunodeficiency virus (HIV)/*Leishmania* coinfecting patients

Coinfecting group	Gender	Age (year)	CD4 <sup>+</sup> T-cell count (cell/mm <sup>3</sup> )	Viral load (log copies/mL)	ART use	Leishmaniasis form
TL/HIV 03	F	42	93	5.4	Unknown	Mucocutaneous active
TL/HIV 05	F	22	291	1.9	Unknown	Cutaneous active
TL/HIV 06	M	36	165	5.3	Unknown	Cutaneous treated
TL/HIV 08	F	62	415	2.7	Yes	Disseminated cutaneous treated
TL/HIV 09	M	59	78	5.0	Yes	Cutaneous active
TL/HIV 12	M	36	615	1.9	Yes	Cutaneous active
TL/HIV 13	M	36	104	5.0	No	Cutaneous active treated
TL/HIV 14	M	25	241	4.6	Yes	Mucocutaneous active

ART: antiretroviral treatment; F: female; M: male; TL: tegumentary leishmaniasis.

CD8<sup>+</sup> T-cell: 0.2, IQR 0.15-0.26) ( $p < 0.05$ ) (Figure). A lymphoproliferative response to SLA was detected in only one coinfecting patient (TL/HIV 06) (12%) who had a previously treated cutaneous leishmaniasis infection. In contrast, CD4<sup>+</sup> and CD8<sup>+</sup> T-cell subsets from all *Leishmania* monoinfected patients proliferated in response to SLA. The CD4<sup>+</sup> and CD8<sup>+</sup> T-cell subsets from both HIV/*Leishmania* coinfecting and *Leishmania* monoinfected patients all proliferated in response to PHA.


The proportions of naïve, CM and EM CD4<sup>+</sup> T-cells obtained after stimulation with SLA were similar between the HIV/*Leishmania* coinfecting and *Leishmania* monoinfected groups (Table II). However, the median absolute counts of naïve, CM and EM CD4<sup>+</sup> T-cells from HIV/*Leishmania* coinfecting patients were significantly lower compared with those of *Leishmania* monoinfected patients ( $p < 0.05$ ). Following SLA stimulation, a statistically significant increase in the median absolute counts of EM CD4<sup>+</sup> T-cells was observed in the cultures from *Leishmania* monoinfected patients ( $p = 0.005$ ). In contrast, there was no change in the median absolute counts of EM CD4<sup>+</sup> T-cells of HIV/*Leishmania* coinfecting patients in the presence of SLA and a statistically significant decrease in the CM CD4<sup>+</sup> T-cell count was observed after culture with SLA ( $p = 0.02$ ). However, there was a statistically significant increase in the absolute number of EM CD4<sup>+</sup> T-cells from both HIV/*Leishmania* coinfecting and *Leishmania* monoinfected groups following PHA stimulation.

**DISCUSSION**

This study demonstrates that a reduction in the lymphoproliferative response to *Leishmania* antigens coincides with a decrease in the absolute numbers of both EM and CM CD4<sup>+</sup> T-cells in response to *Leishmania*

antigens in patients coinfecting with HIV/*Leishmania*. Although the proportion of memory CD4<sup>+</sup> T-cells subsets was similar between monoinfected *Leishmania* and coinfecting HIV/*Leishmania* patients, the median absolute count of EM lymphocytes in coinfecting patients was four-fold lower compared with that of *Leishmania* monoinfected patients. Moreover, following *Leishmania* antigen stimulus, the EM CD4<sup>+</sup> T-cell count of HIV/*Leishmania* coinfecting patients remained below 10 cells/mm<sup>3</sup>, whereas we observed a three-fold increase in the EM CD4<sup>+</sup> T-cell count in the *Leishmania* monoinfected patients. The low number of EM CD4<sup>+</sup> T-cells in HIV/*Leishmania* coinfecting patients might explain the reduced IFN- $\gamma$  production in response to *Leishmania* antigens observed in coinfecting subjects as well as the atypical parasitic spread, frequent relapse and treatment failure observed in these patients (Da-Cruz et al. 1992, Cruz et al. 2006, Lindoso et al. 2009). Although memory and effector CD8<sup>+</sup> T-cell profiles were not evaluated in our study, CD8<sup>+</sup> T-cells from all HIV/*Leishmania* coinfecting patients were unable to proliferate in response to SLA. Previous work has demonstrated the protective role of CD8<sup>+</sup> T-cells in *Leishmania* infection, with healing of lesions associated with high IFN- $\gamma$  production and an increase in the number of CD8<sup>+</sup> T-cells (Da-Cruz et al. 1994, Muller et al. 1994).

The absence of a lymphoproliferative response to *Leishmania* antigens in both the CD4<sup>+</sup> and CD8<sup>+</sup> T-cell subsets examined in our study suggests impairment of the cellular immune response directed at the parasitic infection. Although the HIV/*Leishmania* coinfecting group was heterogeneous with respect to clinical presentation of leishmaniasis (cutaneous vs. mucosal) and stage of treatment (active vs. treated), a significant lack of immune response to *Leishmania* antigens was observed



Proliferative response to *Leishmania* antigens of CD4<sup>+</sup> and CD8<sup>+</sup> T-cells using flow cytometry from human immunodeficiency virus (HIV)/*Leishmania* coinfecting patients with active tegumentary leishmaniasis (TL) (●) and *Leishmania* monoinfected patients (■). Open circles (○) indicate HIV/*Leishmania* coinfecting patients with treated TL. Medians are indicated by horizontal bars. Mann Whitney U test was used to compare groups. Significant differences were found between groups ( $p < 0.05$ ). A: cell division index of CD4<sup>+</sup> T-cells. The dashed line represents the threshold to define positive proliferative response (cell division index > 0.06); B: cell division index of CD8<sup>+</sup> T-cells. The dashed line represents the threshold to define positive proliferative response (cell division index > 0.09). Median cell division index without stimulus (medium): HIV/*Leishmania* co-infected patients 0.02 [interquartile range (IQR) 0.01-0.03] and 0.02 (IQR 0.0-0.04), *Leishmania* monoinfected patients 0.03 (IQR 0.0-0.07) for CD4<sup>+</sup> and CD8<sup>+</sup> T-cell subsets, respectively. Median cell division index following phytohaemagglutinin stimulation: HIV/*Leishmania* co-infected patients 0.7 (IQR 0.2-1.3) and 1 (IQR 0.6-1.4), *Leishmania* only infected patients 0.3 (IQR 0.1-0.5) and 0.2 (IQR 0.1-0.4) for CD4<sup>+</sup> and CD8<sup>+</sup> T-cell subsets, respectively.

across this group of subjects. Only one HIV/*Leishmania* coinfecting patient exhibited a proliferative response to *Leishmania* antigens. This patient was successfully treated for TL before his inclusion in the study and his lesions were in remission. However, the two other coinfecting individuals with treated TL did not develop a proliferative response. Additionally, we found that a decrease in the number of CM CD4<sup>+</sup> T-cells occurred following SLA stimulation in all cultures of coinfecting patients and EM CD4<sup>+</sup> T-cell counts of these patients did not change regardless of presentation of leishmaniasis or stage of treatment. Previous work has shown that the restoration of CD4<sup>+</sup> T-cell-specific responses to antigens in HIV-infected patients undergoing antiretroviral treatment (ART) requires a sustained increase of at least 60 CD4<sup>+</sup> T-cells/ $\mu$ L compared with baseline levels and a reduced viral load (Li et al. 1998). In addition to the quantitative immune defect observed in HIV infection, the quality of the immune response (i.e., degree of immune memory impairment) may also be important in the host response to other pathogens such as *Leishmania*.

Studies in both mice and humans have demonstrated multiple CD4<sup>+</sup> T-cell phenotypes and a broad spectrum of functions have been attributed to these cells in different viral infections (Chen et al. 2001, de Rosa et al. 2001, Roman et al. 2002). Memory T-cells are divided into subpopulations based on the expression of CCR7, a chemokine receptor involved in homing to secondary lymphoid organs and CD45RA, a cell surface molecule involved in T-cell activation. T-cells expressing CD45RA and CCR7 are termed naïve, whereas those with downregulated CD45RA are termed CM cells and produce IL-2. Memory T-cells lacking expression of CCR7 and CD45RA are termed EM T-cells and produce IFN- $\gamma$  (Sallusto et al. 1999). The functional and pheno-

typic heterogeneity of memory T-cells are influenced by variations in antigen exposure and persistence. A dominant single IL-2 and/or IL-2/IFN- $\gamma$  CD4 T-cell response is normally associated with clearance of antigen, whereas a dominant single IFN- $\gamma$  CD4 T-cell response is associated with antigen persistence and high antigen levels (Harari et al. 2005). Effector CD4<sup>+</sup> T-cells are lost in the absence of parasites, whereas CM CD4<sup>+</sup> T-cells are maintained. Upon secondary infection, these CM T-cells become effector T-cells and mediate protection (Zaph et al. 2004). In our study, a decrease in the number of CM CD4<sup>+</sup> T-cells was observed in the coinfecting group following stimulation with SLA. It is possible that during coinfection with HIV and *Leishmania*, the immune activation caused by both HIV and *Leishmania* infection induces increased apoptosis of parasite-specific memory CD4<sup>+</sup> T-lymphocytes (Wolday et al. 1999a, b). The decreased numbers of CM cells and EM cells would reduce the ability of the immune system to develop an adequate systemic response to control the *Leishmania* infection. However, the overall hyporesponsiveness of the cells to *Leishmania* antigen stimulation could also suggest that a qualitative defect may be contributing to this phenomenon. Thus, although an impaired memory response is known to be related to the depletion of CD4<sup>+</sup> T-cells *via* mechanisms related directly or indirectly to HIV (systemic activation and replicative senescence), a defective functional memory response may also be playing a role (Moro-Garcia et al. 2012). Interestingly however, we found that our coinfecting cells responded well to PHA, suggesting that the immunodeficiency was parasite-specific.

Viral suppression by ART may be a contributing factor and in this study, several of our subjects were not receiving ART. In the future, we will examine how the

TABLE II

Quantification of naïve, central (CM) and effector (EM) memory CD4<sup>+</sup> T-cell subsets in response to soluble *Leishmania* antigens (SLA) and to phytohaemagglutinin (PHA) from of human immunodeficiency virus (HIV)/*Leishmania* coinfecting and *Leishmania* monoinfecting groups

CD4 <sup>+</sup> T-cell subset	Medium			SLA			PHA		
	Coinfecting	Monoinfecting	p	Coinfecting	Monoinfecting	p	Coinfecting	Monoinfecting	p
Naïve									
n (%)	30 (17-37)	43 (18-80)	0.8	39 (17-64)	34 (15-57)	0.7	2 (1-4) <sup>b</sup>	4 (2-9) <sup>b</sup>	0.5
n (cell/mm <sup>3</sup> )	9 (3-14)	17 (5-65)	0.02	3 (2-11)	42 (25-106)	0.03	2 (0-6) <sup>b</sup>	13 (4-28)	0.02
CM									
n (%)	23 (8-31)	24 (6-32)	0.6	11 (4-25)	8 (2-23) <sup>a</sup>	0.9	1 (0-4) <sup>b</sup>	2 (1-3)	0.3
n (cell/mm <sup>3</sup> )	55 (31-105)	49 (19-103)	0.001	3 (1-5) <sup>a</sup>	46 (25-119)	0.8	1 (0-3) <sup>b</sup>	8 (2-16) <sup>b</sup>	0.02
EM									
n (%)	12 (11-29)	9 (6-15)	0.2	21 (11-43)	17 (14-23) <sup>a</sup>	0.07	35 (20-40) <sup>b</sup>	30 (25-38) <sup>b</sup>	0.8
n (cell/mm <sup>3</sup> )	8 (5-11)	36 (12-63)	0.0009	6 (2-9)	122 (69-143) <sup>a</sup>	0.0001	27 (11-58) <sup>b</sup>	175 (105-236) <sup>b</sup>	0.0005

Wilcoxon test was used to compare cells cultured with medium and SLA (a) and cells cultured with medium and PHA (b) for each group. Data are presented as median (interquartile range). p: Mann-Whitney U test was used to compare HIV/*Leishmania* coinfecting and *Leishmaniasis* monoinfecting groups.

immune response to *Leishmania* is reconstituted under ART. Although the impact of HIV-1 on host cellular immune function clearly plays an important role in facilitating *Leishmania* infection in dually infected individuals, there is increasing evidence that other mechanisms for the interaction of these two pathogens might also be involved. Particularly, the immunological mechanisms involved in the pathogenesis of mucocutaneous lesions in HIV-infected individuals remain unclear. We hypothesise that innate immunity to *Leishmania* is also compromised in coinfecting patients, resulting in the more severe pathogenesis of mucocutaneous lesions observed in coinfecting patients. Future work will be required to evaluate the effect of coinfection on innate immunity.

#### ACKNOWLEDGEMENTS

To Geraldo Gileno, LPBI, Fiocruz-BA, for providing SLA, and to Maria Zilma Andrade Rodrigues, for your collaboration.

#### REFERENCES

- Bacellar O, D'Oliveira Jr A, Jeronimo S, Carvalho EM 2000. IL-10 and IL-12 are the main regulatory cytokines in visceral leishmaniasis. *Cytokine* 12: 1228-1231.
- Badaro R 1997. When *Leishmania* and HIV interact, a new broad spectrum of leishmaniasis occurs. *Braz J Infect Dis* 1: 145-148.
- Baleeiro CO, Paranhos-Silva M, dos Santos JC, Oliveira GG, Nascimento EG, de Carvalho LP, dos Santos WL 2006. Montenegro's skin reactions and antibodies against different *Leishmania* species in dogs from a visceral leishmaniasis endemic area. *Vet Parasitol* 139: 21-28.
- Bourreau E, Gardon J, Pradinaud R, Pascalis H, Prevot-Linguet G, Karimnia A, Pascal L 2003. Th2 responses predominate during the early phases of infection in patients with localized cutaneous leishmaniasis and precede the development of Th1 responses. *Infect Immun* 71: 2244-2246.
- Carranza-Tamayo CO, de Assis TS, Neri AT, Cupolillo E, Rabello A, Romero GA 2009. Prevalence of *Leishmania* infection in adult HIV/AIDS patients treated in a tertiary-level care center in Brasilia, Federal District, Brazil. *Trans R Soc Trop Med Hyg* 103: 743-748.
- Chen G, Shankar P, Lange C, Valdez H, Skolnik PR, Wu L, Manjunath N, Lieberman J 2001. CD8 T cells specific for human immunodeficiency virus, Epstein-Barr virus and cytomegalovirus lack molecules for homing to lymphoid sites of infection. *Blood* 98: 156-164.
- Coura JR, Galvão-Castro B, Grimaldi Jr G 1987. Disseminated American cutaneous leishmaniasis in a patient with AIDS. *Mem Inst Oswaldo Cruz* 82: 581-582.
- Cruz I, Nieto J, Moreno J, Canavate C, Desjeux P, Alvar J 2006. *Leishmania*/HIV co-infections in the second decade. *Indian J Med Res* 123: 357-388.
- Da-Cruz AM, Conceicao-Silva F, Bertho AL, Coutinho SG 1994. *Leishmania*-reactive CD4<sup>+</sup> and CD8<sup>+</sup> T cells associated with cure of human cutaneous leishmaniasis. *Infect Immun* 62: 2614-2618.
- Da-Cruz AM, Filgueiras DV, Coutinho Z, Mayrink W, Grimaldi Jr G, De Luca PM, Mendonça SCF, Coutinho SG 1999. Atypical mucocutaneous leishmaniasis caused by *Leishmania braziliensis* in an acquired immunodeficiency syndrome patient: T-cell responses and remission of lesions associated with antigen immunotherapy. *Mem Inst Oswaldo Cruz* 94: 537-542.
- Da-Cruz AM, Machado ES, Menezes JA, Rutowitsch MS, Coutinho SG 1992. Cellular and humoral immune responses of a patient with American cutaneous leishmaniasis and AIDS. *Trans R Soc Trop Med Hyg* 86: 511-512.
- de Rosa SC, Herzenberg LA, Roederer M 2001. 11-color, 13-parameter flow cytometry: identification of human naïve T cells by phenotype, function and T-cell receptor diversity. *Nat Med* 7: 245-248.
- UNAIDS - Joint United Nations Program on HIV/AIDS 2012. *Global Report 2012. UNAIDS report on the global AIDS epidemic*, WHO, Geneva, 107 pp.
- Harari A, Vallelian F, Meylan PR, Pantaleo G 2005. Functional heterogeneity of memory CD4 T cell responses in different conditions of antigen exposure and persistence. *J Immunol* 174: 1037-1045.
- Karp CL, Auwaerter PG 2007. Coinfection with HIV and tropical infectious diseases. I. Protozoal pathogens. *Clin Infect Dis* 45: 1208-1213.
- Li TS, Tubiana R, Katlama C, Calvez V, Mohand HA, Autran B 1998. Long-lasting recovery in CD4 T-cell function and viral-load reduction after highly active antiretroviral therapy in advanced HIV-1 disease. *Lancet* 351: 1682-1686.
- Lindoso JA, Barbosa RN, Posada-Vergara MP, Duarte MI, Oyafuso LK, Amato VS, Goto H 2009. Unusual manifestations of tegumentary leishmaniasis in AIDS patients from the New World. *Br J Dermatol* 160: 311-318.
- Machado ES, Braga MP, Da-Cruz AM, Coutinho SG, Vieira ARM, Rutowitsch MS, Cuzzi-Maya T, Grimaldi Jr G, Menezes JA 1992. Disseminated American mucocutaneous leishmaniasis caused by *Leishmania braziliensis braziliensis* in a patient with AIDS: a case report. *Mem Inst Oswaldo Cruz* 87: 487-492.
- Mathur P, Samantaray JC, Vajpayee M, Samanta P 2006. Visceral leishmaniasis/human immunodeficiency virus co-infection in India: the focus of two epidemics. *J Med Microbiol* 55: 919-922.
- Moro-Garcia MA, Alonso-Arias R, Lopez-Larrea C 2012. Molecular mechanisms involved in the aging of the T-cell immune response. *Curr Genomics* 13: 589-602.
- Muller I, Kropf P, Louis JA, Milon G 1994. Expansion of gamma interferon-producing CD8<sup>+</sup> T cells following secondary infection of mice immune to *Leishmania major*. *Infect Immun* 62: 2575-2581.
- Nigro L, Cacopardo B, Preiser W, Braner J, Cinatl J, Palermo F, Russo R, Doerr HW, Nunnari A 1999. In vitro production of type 1 and type 2 cytokines by peripheral blood mononuclear cells from subjects coinfecting with human immunodeficiency virus and *Leishmania infantum*. *Am J Trop Med Hyg* 60: 142-145.
- Pinto LA, Castro BG, Soares MBP, Grassi MFR 2011. An evaluation of the spontaneous proliferation of peripheral blood mononuclear cells in HTLV-1-infected individuals using flow cytometry. *ISRN Oncol* 2011: 326719.
- Rabello A, Orsini M, Disch J 2003. *Leishmania*/HIV co-infection in Brazil: an appraisal. *Ann Trop Med Parasitol* 97 (Suppl. 1): 17-28.
- Rodrigues MZ, Grassi MF, Mehta S, Zhang XQ, Gois LL, Schooley RT, Badaro R 2011. Th1/Th2 cytokine profile in patients coinfecting with HIV and *Leishmania* in Brazil. *Clin Vaccine Immunol* 18: 1765-1769.
- Roman E, Miller E, Harmsen A, Wiley J, von Andrian UH, Huston G, Swain SL 2002. CD4 effector T cell subsets in the response to influenza: heterogeneity, migration and function. *J Exp Med* 196: 957-968.
- Sallusto F, Lenig D, Forster R, Lipp M, Lanzavecchia A 1999. Two subsets of memory T lymphocytes with distinct homing potentials and effector functions. *Nature* 401: 708-712.
- ter Horst R, Collin SM, Ritmeijer K, Bogale A, Davidson RN 2008. Concordant HIV infection and visceral leishmaniasis in Ethiopia:

- the influence of antiretroviral treatment and other factors on outcome. *Clin Infect Dis* 46: 1702-1709.
- Wolday D, Akuffo H, Demissie A, Britton S 1999a. Role of *Leishmania donovani* and its lipophosphoglycan in CD4<sup>+</sup> T-cell activation-induced human immunodeficiency virus replication. *Infect Immun* 67: 5258-5264.
- Wolday D, Berhe N, Akuffo H, Britton S 1999b. *Leishmania*-HIV interaction: immunopathogenic mechanisms. *Parasitol Today* 15: 182-187.
- Wolday D, Berhe N, Akuffo H, Desjeux P, Britton S 2001. Emerging *Leishmania*/HIV co-infection in Africa. *Med Microbiol Immunol* 190: 65-67.
- Wolday D, Berhe N, Britton S, Akuffo H 2000. HIV-1 alters T helper cytokines, interleukin-12 and interleukin-18 responses to the protozoan parasite *Leishmania donovani*. *AIDS* 14: 921-929.
- Zaph C, Uzonna J, Beverley SM, Scott P 2004. Central memory T cells mediate long-term immunity to *Leishmania major* in the absence of persistent parasites. *Nat Med* 10: 1104-1110.